# HAWAIIAN MONK SEAL (Neomonachus schauinslandi)

#### STOCK DEFINITION AND GEOGRAPHIC RANGE

Hawaiian monk seals are distributed throughout the Northwestern Hawaiian Islands (NWHI), with subpopulations at French Frigate Shoals, Laysan Island, Lisianski Island, Pearl and Hermes Reef, Midway Atoll, Kure Atoll, and Necker and Nihoa Islands. They also occur throughout the main Hawaiian Islands (MHI). Genetic variation among monk seals is extremely low and may reflect a long-term history at low population levels and more recent human influences (Kretzmann *et al.* 1997, 2001, Schultz *et al.* 2009). Though monk seal subpopulations often exhibit asynchronous variation in demographic parameters (such as abundance trends and survival rates), they are connected by animal movement throughout the species' range (Johanos *et al.* 2013). Genetic analysis (Schultz *et al.* 2011) indicates the species is a single panmictic population. The Hawaiian monk seal is therefore considered a single stock. Scheel *et al.* (2014) established a new genus, *Neomonachus*, comprising the Caribbean and Hawaiian monk seals, based upon molecular and skull morphology evidence.

#### POPULATION SIZE

The best estimate of the total population size is 1,437 (95% confidence interval 1,367- 1,549; CV = 0.03), (Table 1, Johanos 2019a,b,c). In 2016, new approaches were developed to estimate Hawaiian monk seal abundance, both range-wide and at individual subpopulations (Baker et al. 2016, Harting et al. 2017). In brief, methods for abundance estimation vary by site and year depending on the type and quantity of data available. Total enumeration is the favored method, but requires sufficient field presence to convincingly identify all the seals present, which is typically not achieved at most sites (Baker et al. 2006). When total enumeration is not possible, capture-recapture estimates (using Program CAPTURE) are conducted (Baker 2004; Otis et al. 1978, Rexstad & Burnham 1991, White et al. 1982). When no reliable estimator is obtainable in Program CAPTURE (i.e., the model selection criterion is < 0.75, following Otis et al. 1978), total non-pup abundance is estimated using pre-existing information on the relationship between proportion of the population identified and field effort hours expended (referred to as discovery curve analysis). At rarely visited sites (Necker, Nihoa, Niihau and Lehua Islands) where data are insufficient to use any of the above methods, beach counts are corrected for the proportion of seals at sea. In the MHI other than Niihau and Lehua Islands, abundance is estimated as the minimum tally of all individuals identified by an established sighting network during the calendar year. At all sites, pups are tallied. Finally, site-specific abundance estimates and their uncertainty are combined using Monte Carlo methods to obtain a range-wide abundance estimate distribution. All the above methods are described or referenced in Baker et al. (2016) and Harting et al. (2017). Note that because some of the abundance estimation methods utilize empirical distributions which are updated as new data accrue, previous years' estimates can change slightly when recalculated using these updated distributions.

In 2018, total enumeration was achieved at Laysan and Lisianski Islands, and Kure Atoll, and a capture-recapture estimate was obtained for Midway Atoll. At French Frigate Shoals and Pearl and Hermes Reef, abundance estimates were obtained using discovery curve analysis (Table 1). Counts at Necker and Nihoa Islands are conducted from zero to a few times per year. Pups are born over the course of many months and have very different haulout patterns compared to older animals. Therefore, pup production at Necker and Nihoa Islands is estimated as the mean of the total pups observed in the past 5 years, excluding counts occurring early in the pupping season when most have vet to be born.

In the MHI, NMFS collects information on seal sightings reported throughout the year by a variety of sources, including a volunteer network, the public, and directed NMFS observation effort. In recent years, a small number of surveys of Ni'ihau and nearby Lehua Islands have been conducted through a collaboration between NMFS, Ni'ihau residents and the US Navy. Total MHI monk seal abundance is estimated by adding the number of individually identifiable seals documented during a calendar year on all MHI other than Ni'ihau and Lehua to an estimate for these latter two islands based on counts expanded by a haulout correction factor. A recent telemetry study (Wilson *et al.*, 2017) found that MHI monk seals (N=23) spent a greater proportion of time ashore than Harting *et al.* (2017) estimated for NWHI seals. Therefore, the total non-pup estimate for Ni'ihau and Lehua Islands was the total beach count at those sites (less individual seals already counted at other MHI) divided by the mean proportion of time hauled out in the MHI (Wilson *et al.*, 2017). The total pups observed at Ni'ihau and Lehua Islands were added to obtain the total (Table 1).

Table 1. Total and minimum estimated abundance ( $N_{min}$ ) of Hawaiian monk seals by location in 2018. The estimation method is indicated for each site. Methods used include DC: discovery curve analysis, EN: total enumeration; CR: capture-recapture; CC: counts corrected for the proportion of seals at sea; Min: minimum tally. Median values are presented. Note that the median range-wide abundance is not equal to the total of the individual sites' medians, because the median of sums may differ from the sum of medians for non-symmetrical distributions.  $N_{min}$  for individual sites are either the minimum number of individuals identified or the  $20^{th}$  percentile of the abundance distribution (the latter applies to Necker, Nihoa, Ni'ihau/Lehua, and range-wide).

	Total			Nmin			
Location	Non-pups	Pups	Total	Non-pups	Pups	Total	Method
French Frigate Shoals	181	40	221	181	40	221	DC
Laysan	200	30	230	200	30	230	EN
Lisianski	130	15	145	130	15	145	EN
Pearl and Hermes Reef	124	26	150	124	26	150	DC
Midway	71	13	84	65	13	78	CR
Kure	91	11	102	91	11	102	EN
Necker	76	7	83	64	7	71	CC
Nihoa	100	6	106	85	6	91	CC
MHI_(without Ni'ihau/ Lehua)	145	30	175	145	30	175	Min
Ni'ihau/Lehua	124	7	131	104	7	111	CC
Range-wide	1252	185	1437	1189	185	1374	

## **Minimum Population Estimate**


The total numbers of seals identified at the NWHI subpopulations other than Necker and Nihoa, and in the MHI other than Ni'ihau and Lehua, are the best estimates of minimum population size at those sites. Minimum population sizes for Necker, Nihoa, Ni'ihau, and Lehua Islands are estimated as the lower 20<sup>th</sup> percentiles of the non-pup abundance distributions generated using haulout corrections as described above, plus the pup estimates. The minimum abundance estimates for each site and for all sites combined (1,374) are presented in Table 1.

### **Current Population Trend**

Range-wide abundance estimates are available from 2013 to 2018 (Figure 1). While these estimates remain somewhat negatively-biased for reasons explained in Baker et al. (2016), they provided a much more comprehensive assessment of status and trends than has been previously available. A Monte Carlo approximation of the annual multiplicative rate of realized population growth during 2013-2018 was generated by fitting 10,000 log-linear regressions to randomly selected values from each year's abundance distributions. The median rate (and 95% confidence limits) is 1.02 (1.00, 1.04). Thus, the best estimate is that the population grew at an average rate of about 2% per year from 2013 to 2018. Less than 1% of the distribution was below 1, indicating that there is greater than a 99% chance that the monk seal population increased during 2013-2018.

## **CURRENT AND MAXIMUM NET PRODUCTIVITY RATES**

Mean non-pup beach counts are used as a long-term index of abundance for years when data are insufficient to estimate total abundance as described above. Prior to 1999, beach count increases of up to 7% annually were observed at Pearl and Hermes Reef, and this is the highest estimate of the maximum net productivity rate ( $R_{max}$ ) observed for this species (Johanos 2019a). Consistent with this value, a life table analysis representing a time when the MHI monk seal population was apparently expanding, yielded an estimated intrinsic population growth rate of 1.07 (Baker *et al.* 2011).



**Figure 1**. Range-wide abundance of Hawaiian monk seals, 2013-2018. Medians and 95% confidence limits are shown. Estimates prior to 2018 are re-estimated based on new data and represent negligible changes compared with values reported in the final 2019 stock assessment.

#### POTENTIAL BIOLOGICAL REMOVAL

Using current minimum population size (1,374),  $R_{max}$  (0.07) and a recovery factor  $(F_r)$  for ESA endangered stocks (0.1), yields a Potential Biological Removal (PBR) of 4.8.

#### **HUMAN-CAUSED MORTALITY AND SERIOUS INJURY**

Human-related mortality has caused two major declines of the Hawaiian monk seal (Ragen 1999). In the 1800s, this species was decimated by sealers, crews of wrecked vessels, and guano and feather hunters (Dill and Bryan 1912; Wetmore 1925; Bailey 1952; Clapp and Woodward 1972). Following a period of at least partial recovery in the first half of the 20<sup>th</sup> century (Rice 1960), most subpopulations again declined. This second decline has not been fully explained, but long-term trends at several sites appear to have been driven both by variable oceanic productivity (represented by the Pacific Decadal Oscillation) and by human disturbance (Baker *et al.* 2012, Ragen 1999, Kenyon 1972, Gerrodette and Gilmartin 1990). Currently, human activities in the NWHI are limited and human disturbance is relatively rare, but human-seal interactions, have become an important issue in the MHI. Intentional killing of seals in the MHI is an ongoing and serious concern (Table 2).

**Table 2**. Intentional and potentially intentional killings of MHI monk seals, and anthropogenic mortalities not associated with fishing gear during 2014-2018 (Johanos 2019d, Mercer 2020). There were no confirmed cases in 2016.

Year	Age/sex	Island	Cause of Death	Comments
2014	Adult male	Oahu	Suspected trauma	Intent unconfirmed
2014	Pup female	Kauai	Skull fracture, blunt force trauma	Likely intentional
2015	Pup male	Kauai	Dog attack/bite wounds	4 other seals injured during this event
2015	Juvenile male	Kauai	Probable boat strike	
2015	Adult male	Laysan	Research handling	Accidental, specific cause undetermined
2017	Adult female	Kauai	Trauma	Suspect intentional
2017	Juvenile female	Molokai	Blunt force trauma	Suspect intentional
2018	Juvenile female	Molokai	Blunt force trauma	Intentional

It is extremely unlikely that all carcasses of intentionally killed monk seals are discovered and reported. Studies of the recovery rates of carcasses for other marine mammal species have shown that the probability of detecting and documenting most deaths (whether from human or natural causes) is quite low (Peltier *et al.* 2012; Williams *et al.* 2011; Perrin *et al.* 2011; Punt and Wade 2010).

#### **Fishery Information**

Fishery interactions with monk seals can include direct interaction with gear (hooking or entanglement), seal consumption of discarded catch, and competition for prey. Entanglement of monk seals in derelict fishing gear, which is believed to originate outside the Hawaiian archipelago, is described in a separate section. Fishery interactions are a serious concern in the MHI, especially involving nearshore fisheries managed by the State of Hawaii (Gobush et al. 2016). There are no fisheries operating in or near the NWHI. In 2018, ten seal hookings were documented, two of which were classified as serious and eight as non-serious injuries. Of the non-serious injuries, two would have been deemed serious had they not been mitigated (Henderson 2019a, Mercer 2020). The hooks involved included circle, treble and J-hooks of widely varying sizes. Monk seals also interact with nearshore gillnets, and several confirmed deaths have resulted. One confirmed gillnet mortality occurred in 2017, and three mortalities in 2016-2017 are considered suspect net mortalities (Mercer 2018), based on necropsy findings of probable peracute underwater entrapment (drowning) (Moore et al. 2013). In 2018, a fisherman reported releasing a monk seal from a gillnet he was tending off Oahu. The seal was reportedly lethargic but the event was deemed non-serious because the seal was released and subsequently has been resighted alive. Two seals were entangled in monofilament fishing gear on Oahu in 2018; both were deemed non-serious because they were disentangled, but would have been serious had they not been mitigated. No mortality or serious injuries have been attributed to the MHI bottomfish handline fishery (Table 3). Published studies on monk seal prey selection based upon scat/spew analysis and video from seal-mounted cameras revealed evidence that monk seals fed on families of bottomfish which contain commercial species (many prey items recovered from scats and spews were identified only to the level of family; Goodman-Lowe 1998, Longenecker et al. 2006, Parrish et al. 2000). Quantitative fatty acid signature analysis (QFASA) results support previous studies illustrating that monk seals consume a wide range of species (Iverson et al. 2011). However, deepwater-slope species, including two commercially targeted bottomfishes and other species not caught in the fishery, were estimated to comprise a large portion of the diet for some individuals. Similar species were estimated to be consumed by seals regardless of location, age or gender, but the relative importance of each species varied. Diets differed considerably between individual seals. These results highlight the need to better understand potential ecological interactions with the MHI bottomfish handline fishery.

**Table 3.** Summary of mortality, serious and non-serious injury of Hawaiian monk seals due to fisheries and calculation of annual mortality rate. n/a indicates that sufficient data are not available. Percent observer coverage for the deep and shallow-set components, respectively, of the pelagic longline fishery, are shown. Total non-serious injuries are presented as well as, in parentheses, the number of those injuries that would have been deemed serious had they not been mitigated (e.g., by de-hooking or disentangling). Data for MHI bottomfish and nearshore fisheries are based upon incidental observations (i.e., hooked seals and those entangled in active gear). All hookings not clearly attributable to either fishery with certainty were attributed to the bottomfish fishery, and hookings which resulted in injury of unknown severity were classified as serious. Nearshore fisheries injuries and mortalities include seals entangled/drowned in nearshore gillnets and hooked/entangled in hook-and-line gear, recognizing that it is not possible to determine whether the nets or hook-and-line gear involved were being used for commercial purposes.

Fishery Name	Year	Data Type	% Obs. coverage	Observed/Reported Mortality/Serious Injury	Estimated Mortality/ Serious Injury	Non-serious (Mitigated serious)	Mean Takes (CV)
Pelagic Longline	2014 2015 2016 2017 2018	observer observer observer observer observer	20.8% & 100% 20.6% & 100% 20.1% & 100% 20.4% & 100% 20.4% & 100%	0 0 0 0 0	0 0 0 0	0 0 0 0	0 (0)
MHI Bottomfish			none		n/a		n/a

	2014 2015 2016 2017 2018	Incidental observations of seals		0 0 0 0 0		0 0 0 0	
Nearshore	2014 2015 2016 2017 2018	Incidental observations of seals	none	1 3 0 3 2	n/a	13 (9) 8 (2) 11 (6) 19(6) 11(4)	≥1.8
Mariculture	2014 2015 2016 2017 2018	Incidental Observation	none	0 0 0 1 0	n/a	0 0 0 0	0.2 (2.2)
Minimum total annual takes							≥ 2.0

### **Fishery Mortality Rate**

Total fishery mortality and serious injury is not considered to be insignificant and approaching a rate of zero. Monk seals are being hooked and entangled in the MHI at a rate that has not been reliably assessed but is certainly greater than zero. The information above represents only reported direct interactions, and without directed observation effort, the true interaction rate cannot be estimated. Monk seals also die from entanglement in fishing gear and other debris throughout their range (likely originating from various sources outside of Hawaii), and NMFS along with partner agencies is pursuing a program to mitigate entanglement (see below).

#### **Entanglement in Marine Debris**

Hawaiian monk seals become entangled in fishing and other marine debris at rates higher than reported for other pinnipeds (Henderson 2001). Several hundred cases of debris entanglement have been documented in monk seals (nearly all in the NWHI), including ten documented deaths (one of which occurred at Kure Atoll in 2018) (Henderson 2001; Henderson 2019b, Mercer 2020). The number of marine debris entanglements documented in the past five years (Table 4) is an underestimate of the total impact of this threat because no people are present to document nor mitigate entanglements at most of the NWHI for the majority of the year. Nearly all documented cases would have been deemed serious had they not been mitigated by field biologists. The fishing gear fouling the reefs and beaches of the NWHI and entangling monk seals only rarely includes types used in Hawaii fisheries. For example, trawl net and monofilament gillnet accounted for approximately 35% and 34%, respectively, of the debris removed from reefs in the NWHI by weight, and trawl net alone accounted for 88% of the debris by frequency (Donohue *et al.* 2001), despite the fact that trawl fisheries have been prohibited in Hawaii since the 1980s.

**Table 4.** Summary of documented marine debris entanglements of Hawaiian monk seals during the most recent five years. Total non-serious injuries are presented as well as, in parentheses, the number of those injuries that would have been deemed serious had the seals not been disentangled.

Year	Observed/Reported Mortality/Serious Injury	Non-serious (Mitigated serious)
2014	0	5(4)
2015	0	12(8)
2016	0	3(2)
2017	0	11(8)
2018	1	13(4)
Minimum total annual takes	≥ 0.2	

The NMFS and partner agencies continue to mitigate impacts of marine debris on monk seals as well as turtles, coral reefs and other wildlife. Marine debris is removed from beaches and seals are disentangled during annual population assessment activities at the main reproductive sites. Since 1996, annual debris survey and removal efforts in the NWHI coral reef habitat have been ongoing (Donohue *et al.* 2000, Donohue *et al.* 2001, Dameron *et al.* 2007).

#### **Toxoplasmosis**

Land-to-sea transfer of *Toxoplasma gondii*, a protozoal parasite shed in the feces of cats, is of growing concern. Although the parasite can infect many species, felids are the definitive host, meaning it can only reproduce in cats. There are no native felids in Hawaii, but several hundred thousand feral and domestic cats occur throughout the MHI. As such, all monk seal deaths attributable to toxoplasmosis are considered human caused. A case definition for toxoplasmosis and other protozoal-related mortalities was developed and retrospectively applied to 306 cases of monk seal mortality from 1982-2015 (Barbieri et al. 2016). During the past five years (2014-2018) five monk seal deaths (representing a minimum average of one death per year) have been directly attributed to toxoplasmosis (Mercer 2020). All five deaths involved female seals. The number of deaths from this pathogen are likely underrepresented, given that more seals disappear each year than are found dead and examined, and the potential for chronic infections remains poorly understood in this species. Furthermore, *T. gondii* can be transmitted vertically from dam to fetus, and failed pregnancies are difficult to detect in wild, free-ranging animals. Unlike threats such as hook ingestion or malnutrition, which can often be mitigated through rehabilitation, options for treating seals with toxoplasmosis are severely restricted. The accumulating number of monk seal deaths from toxoplasmosis in recent years is a growing concern given the increasing geographic overlap between humans, cats, and Hawaiian monk seals in the MHI.

#### **Other Mortality**

Sources of mortality that impede recovery include food limitation (see Habitat Issues), single and multiplemale intra-species aggression (mobbing), shark predation, and disease/parasitism. Male seal aggression has caused episodes of mortality and injury. Past interventions to remove aggressive males greatly mitigated, but have not eliminated, this source of mortality (Johanos *et al.* 2010). Galapagos shark predation on monk seal pups has been a chronic and significant source of mortality at French Frigate Shoals since the late 1990s, despite mitigation efforts by NMFS (Gobush 2010). Besides toxoplasmosis, infectious disease effects on monk seal demographic trends are low relative to other stressors. However, a disease outbreak introduced from livestock, feral animals, pets or other carrier wildlife could be catastrophic to the immunologically naïve monk seal population. Key disease threats include West Nile virus, morbillivirus and influenza.

#### **Habitat Issues**

Poor juvenile survival rates and variability in the relationship between weaning size and survival suggest that prey availability has limited recovery of NWHI monk seals (Baker and Thompson 2007, Baker *et al.* 2007, Baker 2008). Multiple strategies for improving juvenile survival, including translocation and captive care are being implemented (Baker and Littnan 2008, Baker *et al.* 2013, Norris 2013). A testament to the effectiveness of past actions to improve survival, Harting *et al.* (2014) demonstrated that approximately one-third of the monk seal population alive in 2012 was made up of seals that either had been intervened with to mitigate life-threatening situations, or were descendants of such seals. In 2014, NMFS produced a final Programmatic Environmental Impact Statement (PEIS) on current and future anticipated research and enhancement activities and issued a permit covering the activities described in the PEIS preferred alternative. Loss of terrestrial habitat at French Frigate Shoals is a serious threat to the viability of the resident monk seal population. Prior to 2018, pupping and resting islets had shrunk or virtually disappeared (Antonelis *et al.* 2006). In 2018, the two remaining primary islands where pups were born at French Frigate Shoals (Trig and East Islands) were obliterated due to progressive erosion and hurricane Walaka (in September 2018). Projected increases in global average sea level are expected to further significantly reduce terrestrial habitat for monk seals in the NWHI (Baker *et al.* 2006, Reynolds *et al.* 2012).

The seawall at Tern Island, French Frigate Shoals, continues to degrade and poses an increasing entrapment hazard for monk seals and other fauna. The situation has worsened since 2012, when the USFWS ceased operations on Tern Island, thus leaving the island unmanned for most of the year. Previously, daily surveys were conducted throughout the year to remove entrapped animals. Now this only occurs when NMFS monk seal field staff are on site. Furthermore, sea wall breaches are allowing sections of the island to erode and undermine buildings and other infrastructure. Several large water tanks have collapsed, exposing pipes and wiring that may entangle or entrap seals. In September 2018, hurricane Walaka exacerbated this situation by largely destroying remaining structures and strewing the resulting debris around the island. Strategies to mitigate these threats are currently under consideration.

Goodman-Lowe (1998) provided information on prey selection using hard parts in scats and spewings. Information on at-sea movement and diving is available for seals at all six main subpopulations in the NWHI using satellite telemetry (Stewart *et al.* 2006). Cahoon (2011) and Cahoon *et al.* (2013) described diet and foraging behavior of MHI monk seals, and found no striking difference in prey selection between the NWHI and MHI.

Monk seal juvenile survival rates are favorable in the MHI (Baker et al. 2011). Further, the excellent

condition of pups weaned on these islands suggests that there are ample prey resources available, perhaps in part due to fishing pressure that has reduced monk seal competition with large fish predators (sharks and jacks) (Baker and Johanos 2004). Yet, there are many challenges that may limit the potential for growth in this region. The human population in the MHI is approximately 1.4 million compared to fewer than 100 in the NWHI, such that anthropogenic threats in the MHI are considerable. Intentional killing of seals is a very serious concern. Also, the same fishing pressure that may have reduced the monk seal's competitors is a source of injury and mortality. Vessel traffic in the populated islands entails risk of collision with seals and impacts from oil spills. A mortality in 2015 was deemed most likely due to boat strike. Finally, as noted above, toxoplasmosis is now recognized as a serious anthropogenic threat to seals in the MHI.

#### STATUS OF STOCK

In 1976, the Hawaiian monk seal was designated depleted under the Marine Mammal Protection Act of 1972 and as endangered under the Endangered Species Act of 1973. Therefore, the Hawaiian monk seal is a strategic stock. The species is well below its optimum sustainable population and has not recovered from past declines. Annual human-caused mortality for the most recent 5-year period (2014-2018) was at least 4.8 animals, which equals PBR, including fishery-related mortality in nearshore gillnets, hook-and-line gear, and mariculture ( $\geq 2.0$ /yr, Table 3), intentional killings and other human-caused mortalities ( $\geq 1.6$ /yr, Table 2), entanglement in marine debris ( $\geq 0.2$ /yr, Table 4), and deaths due to toxoplasmosis ( $\geq 1.0$ /yr).

#### REFERENCES

- Antonelis, G.A., J.D. Baker, T.C. Johanos, R.C. Braun, and A.L. Harting. 2006. Hawaiian monk seal (*Monachus schauinslandi*): Status and Conservation Issues. Atoll Res. Bull. 543:75-101.
- Bailey, A.M. 1952. The Hawaiian monk seal. Museum Pictorial, Denver Museum of Natural History 7:1-32.
- Baker, J.D. 2004. Evaluation of closed capture-recapture methods to estimate abundance of Hawaiian monk seals, *Monachus schauinslandi*. Ecol. Appl. 14:987-998.
- Baker J.D. 2008. Variation in the relationship between offspring size and survival provides insight into causes of mortality in Hawaiian monk seals. Endangered Species Research 5:55-64.
- Baker J.D., A.L. Harting, and C.L. Littnan. 2013. A two-stage translocation strategy for improving juvenile survival of Hawaiian monk seals. Endang. Species Res. 21:33–44.
- Baker, J.D., A.L. Harting, and T.C. Johanos. 2006. Use of discovery curves to assess abundance of Hawaiian monk seals. Mar. Mamm. Sci. 22:847-861.
- Baker J.D., A.L. Harting, T.C. Johanos, and C.L. Littnan. 2016. Estimating Hawaiian monk seal range-wide abundance and associated uncertainty. Endang. Species Res. 31:317-324.
- Baker J.D., A. L. Harting, T.A. Wurth, and T.C, Johanos. 2011. Dramatic shifts in Hawaiian monk seal distribution predicted from divergent regional trends. Mar. Mamm. Sci. 27: 78–93.
- Baker, J.D. and T.C. Johanos. 2004. Abundance of Hawaiian monk seals in the main Hawaiian Islands. Biol. Conserv. 116:103-110.
- Baker J.D., and Littnan C.L. 2008. Report of the Hawaiian Monk Seal Captive Care Workshop, Honolulu, Hawaii, June 11–13, 2007. Pacific Islands Fish. Sci. Cent., Natl. Mar. Fish. Serv., NOAA, Honolulu, HI 96822-2396. Pacific Islands Fish. Sci. Cent. Admin. Rep. H-08-02, 42 p.
- Baker J.D., E.A. Howell, and J.J. Polovina. 2012. Relative influence of climate variability and direct anthropogenic impact on a sub-tropical Pacific top predator, the Hawaiian monk seal. Mar. Ecol. Prog. Ser. 469:175-189.
- Baker J.D., C.L. Littnan, and D.W. Johnston. 2006. Potential effects of sea-level rise on the terrestrial habitats of endangered and endemic megafauna in the Northwestern Hawaiian Islands. Endang. Species Res. 4:1-10.
- Baker, J.D., J.J. Polovina, and E.A. Howell. 2007. Effect of variable oceanic productivity on the survival of an upper trophic predator, the Hawaiian monk seal, Monachus schauinslandi. Mar. Ecol. Prog. Ser. 346:277-283.
- Baker J.D. and P.M. Thompson. 2007. Temporal and spatial variation in age-specific survival rates of a long-lived mammal, the Hawaiian monk seal. Proc. Roy. Soc. B 274:407-415.
- Barbieri, M.M., L. Kashinsky, D.S. Rotstein, K.M. Colegrove, K.H. Haman, S.L. Magargal, A.R. Sweeny, A.C. Kaufman, M.E. Grigg, and C.L. Littnan. 2016. Protozoal-related mortalities in endangered Hawaiian monk seals Neomonachus schauinslandi. Dis. Aquat. Organisms 121:85-95.
- Cahoon, M.K. 2011. The foraging ecology of monk seals in the main Hawaiian Islands. MSc thesis, University of Hawaii, 172 p.
- Cahoon M.K., C.L. Littnan, K. Longenecker, and J.R. Carpenter. 2013. Dietary comparison of two Hawaiian monk seal populations: the role of diet as a driver of divergent population trends. Endang. Species Res. 20:137-

146.

- Clapp, R.B., and P.W. Woodward. 1972. The natural history of Kure Atoll, Northwestern Hawaiian Islands, Atoll Res. Bull. 164:303-304.
- Dameron OJ., M. Park, M. Albins, and R. Brainard. 2007. Marine debris accumulation in the Northwestern Hawaiian Islands: An examination of rates and processes. Mar. Pollut. Bull. 54(4): 423-433.
- Dill, H.R., and W.A. Bryan. 1912. Report on an expedition to Laysan Island in 1911. U.S. Dept. of Agric. Surv. Bull. 42:1-30.
- Donohue, M.J., R. Brainard, M. Parke, and D. Foley. 2000. Mitigation of environmental impacts of derelict fishing gear through debris removal and environmental monitoring. *In* Hawaiian Islands Humpback Whale National Marine Sanctuary, Proceedings of the International Marine Debris Conference on Derelict Fishing Gear and the Ocean Environment, 6-11 August 2000, Honolulu, Hawaii. p. 383-402.
- Donohue, M.J., R.C. Boland, C.M. Sramek, and G.A. Antonelis. 2001. Derelict fishing gear in the Northwestern Hawaiian Islands: diving surveys and debris removal in 1999 confirm threat to coral reef ecosystems. Mar. Pollut. Bull. 42:1301-1312.
- Gerrodette, T.M., and W.G. Gilmartin. 1990. Demographic consequences of changed pupping and hauling sites of the Hawaiian monk seal. Conserv. Biol. 4:423-430.
- Gobush, K.S. 2010. Shark predation on Hawaiian monk seals: Workshop II & post-workshop developments, November 5-6, 2008. U.S. Dep. Commer., NOAA Tech. Memo., NOAA-TM-NMFS-PIFSC-21, 43 p. + Appendices.
- Gobush, K.S., Wurth, T.A., Henderson, J. R., Becker, B. L., and C.L. Littnan. 2016. <u>Prevalence of interactions between Hawaiian monk seals (Nemonachus schauinslandi)</u> and nearshore fisheries in the main Hawaiian <u>Islands. Pac. Conserv. Biol. 23:25-31.</u>
- Goodman-Lowe, G.D. 1998. Diet of the Hawaiian monk seal (*Monachus schauinslandi*) from the northwestern Hawaiian islands during 1991 to 1994. Mar. Biol. 132:535-546.
- Harting AL, Baker JD, Johanos TC. 2017. Estimating population size for Hawaiian monk seals using haulout data. J. Wildl. Manage. 81:1202-9.
- Harting A.L., T.C. Johanos, and C.L. Littnan. 2014. <u>Benefits derived from opportunistic survival-enhancing interventions for the Hawaiian monk seal: the silver BB paradigm. Endang. Spec. Res. 25: 89-96.</u>
- Henderson, J.R. 1990. Recent entanglements of Hawaiian monk seals in marine debris. <u>In</u> R. S. Shomura and M. L. Godfrey (eds.), Proceedings of the Second International Conference on Marine Debris, April 2-7, 1989, Honolulu, Hawaii, p. 540-553. U.S. Dep. Commer., NOAA, Tech. Memo. NMFS-SWFSC-154.
- Henderson, J.R. 2001. A Pre and Post MARPOL Annex V Summary of Hawaiian Monk Seal Entanglements and Marine Debris Accumulation in the Northwestern Hawaiian Islands, 1982-1998. Mar. Pollut. Bull. 42:584-589
- Henderson, J.R. 2019a. Hawaiian Monk Seal Research Program Hawaiian monk seal fisheries interaction data collected in the Hawaiian Archipelago, 1976-2018. US National Oceanographic Data Center.
- Henderson, J.R. 2019b. Hawaiian Monk Seal Research Program Hawaiian monk seal entanglement data collected in the Hawaiian Archipelago, 1974-2018. US National Oceanographic Data Center.
- Iverson, S., J. Piché, and W. Blanchard. 2011. Hawaiian monk seals and their prey: assessing characteristics of prey species fatty acid signatures and consequences for estimating monk seal diets using Quantitative Fatty Acid Signature Analysis. U.S. Dep. Commer., NOAA Tech. Memo., NOAA-TM-NMFS-PIFSC-23, 114 p. + Appendices.
- Johanos, T.C. 2019a. Hawaiian Monk Seal Research Program Hawaiian monk seal survey data collected in the Hawaiian Archipelago, 1981-2018. US National Oceanographic Data Center.
- Johanos, T.C. 2019b. Hawaiian Monk Seal Research Program Hawaiian monk seal master identification records (annual) collected in the Hawaiian Archipelago, 1981-2018. US National Oceanographic Data Center.
- Johanos, T.C. 2019c. Hawaiian Monk Seal Research Program Hawaiian monk seal master identification records (seal) collected in the Hawaiian Archipelago, 1981-2018. US National Oceanographic Data Center.
- Johanos, T.C. 2019d. Hawaiian Monk Seal Research Program Hawaiian monk seal survival factors collected in the Hawaiian Archipelago, 1977-2018. US National Oceanographic Data Center.
- Johanos, T.C. and J.D. Baker (editors). 2001. The Hawaiian monk seal in the Northwestern Hawaiian Islands, 1999. U.S. Dep. Commer., NOAA Tech. Memo. NOAA-TM-NMFS-SWFSC-310, 130 p.
- Johanos T.C., B.L. Becker, J.D. Baker, T.C. Ragen, W.G. Gilmartin, and T. Gerrodette. 2010. Impacts of sex ratio reduction on male aggression in the critically endangered Hawaiian monk seal *Monachus schauinslandi*. Endang. Species Res. 11:123–132.

- Johanos, T.C., A.L. Harting, T.A. Wurth, and J.D. Baker. 2013. <u>Range-wide movement patterns of Hawaiian monk</u> seals. Mar. Mamm. Sci. 30:1165-1174.
- Kenyon, K.W. 1972. Man versus the monk seal. J. Mammal. 53:687-696.
- Kretzmann, M.B., W.G. Gilmartin, A. Meyer, G.P. Zegers, S.R. Fain, B.F. Taylor, and D.P. Costa. 1997. Low genetic variability in the Hawaiian monk seal. Conserv. Biol. 11:482-490.
- Kretzmann, M.B., N.J. Gemmell, and A. Meyer. 2001. Microsatellite analysis of population structure in the endangered Hawaiian monk seal. Conserv. Biol. 15:457-466.
- Mercer, T.A. 2020. Summary of documented human-caused mortality, serious injury and non-serious injury in Hawaiian monk seals: supporting documentation for the 2020 Stock Assessment Report.
- Moore, M. J., J. van der Hoop, S. G. Barco, A. M. Costidis, F. M. Gulland, P. D. Jepson, K. T. Moore, S. Raverty, and W. A. McLellan. 2013. Criteria and case definitions for serious injury and death of pinnipeds and cetaceans caused by anthropogenic trauma. Diseases of Aquatic Organisms 103: 229-264.
- Norris, T.A. 2013. Foraging behavior, habitat, health, and survival of resident and translocated Hawaiian monk seals at Nihoa Island, Hawaii. Msc thesis, San Jose State University, 121 p.
- Otis, D. L., K.P. Burnham, G.C. White, and D.R. Anderson. 1978. Statistical inference from capture data on closed animal populations. Wildl. Monogr. 62:1-135.
- Punt, A.E., and P.R. Wade. 2010. Population status of the eastern North Pacific stock of gray whales in 2009. U.S. Dep. Commer. NOAA Tech. Memo. NMFS-AFSC-207, 43 p.
- Parrish, F.A., M.P. Craig, T.J. Ragen, G.J. Marshall, and B.M. Buhleier. 2000. Identifying diurnal foraging habitat of endangered Hawaiian monk seals using a seal-mounted video camera. Mar. Mamm. Sci. 16:392-412.
- Peltier, H., W. Dabin, P. Daniel, O. Van Canneyt, G. Dorémus, M. Huon, and V. Ridoux. 2012. The significance of stranding data as indicators of cetacean populations at sea: Modelling the drift of cetacean carcasses. Ecol. Indicators 18:278-290.
- Perrin, W.F., J.L. Thieleking, W.A. Walker, F.I. Archer, and K.M. Robertson. 2011. Common bottlenose dolphins (*Tursiops truncatus*) in California waters: Cranial differentiation of coastal and offshore ecotypes. Mar. Mamm. Sci. 27:769-792.
- Ragen, T.J. 1993. Status of the Hawaiian monk seal in 1992. Admin. Rep. H-93-05. Southwest Fisheries Science Center, National Marine Fisheries Service, 2570 Dole St., Honolulu, HI 96822-2396. 79 pp.
- Ragen, T.J. 1999. Human activities affecting the population trends of the Hawaiian monk seal. Pages 183-194 *in* J.A. Musick, ed. Life in the slow lane: Ecology and conservation of long-lived marine animals. American Fisheries Society Symposium 23, American Fisheries Society, Bethesda, MD.
- Reynolds, M.H., P. Berkowitz, K.N. Courtot, and C.M. Krause, eds. 2012. Predicting sea-level rise vulnerability of terrestrial habitat and wildlife of the Northwestern Hawaiian Islands: U.S. Geological Survey Open-File Report 2012–1182, 139 p.
- Rexstad, E.A., and K.P. Burnham. 1991. User's manual for interactive Program CAPTURE. Colorado Cooperative Fish and Wildlife Research Unit, Colorado State University, Fort Collins, CO. 29 pp.
- Rice, D.W. 1960. Population dynamics of the Hawaiian monk seal. J. Mammal. 41:376-385.
- Scheel, D.M., G.J. Slater, S.O. Kolokotronis, C.W. Potter, D.S. Rotstein, K. Tsangaras, A.D. Greenwood, and K.M. Helgen. 2014. Biogeography and taxonomy of extinct and endangered monk seals illuminated by ancient DNA and skull morphology. ZooKeys 409:1-33.
- Schultz J.K., Baker J.D., Toonen R.J., Bowen B.W. 2009. Extremely low genetic diversity in the endangered Hawaiian monk seal (*Monachus schauinslandi*). J. Heredity 100:25-33.
- Schultz J.K., J.D. Baker, R.J. Toonen, A.L. Harting, and B.W. Bowen. 2011. Range-wide genetic connectivity of the Hawaiian monk seal and implications for translocation. Conserv. Biol. 25:124-132.
- Stewart B.S., G.A. Antonelis, J.D. Baker, and P.Y. Yochem. 2006. Foraging biogeography of the Hawaiian monk seal in the Northwestern Hawaiian Islands. Atoll Res Bull 543:131-145. Wade, P.R. and R.P. Angliss. 1997. Guidelines for Assessing Marine Mammal Stocks: Report of the GAMMS Workshop April 3-5, 1996, Seattle, Washington. U. S. Dep. Commer., NOAA Tech. Memo. NMFS-OPR-12. 93 pp.
- Wade, P.R. 1998. Calculating limits to the allowable human-caused mortality of cetaceans and pinnipeds. Mar. Mamm. Sci. 14:1-37.
- Wetmore, A. 1925. Bird life among lava rock and coral sand. The Natl. Geograp. Mag. 48:77-108.
- White, G.C., D.R. Anderson, K.P. Burnham, and L. Otis. 1982. Capture-recapture and removal methods for sampling closed populations. Los Alamos National Laboratory, Los Alamos, New Mexico.
- Williams, R., S. Gero, L. Bejder, J. Calambokidis, S.D. Kraus, D. Lusseau, A.J. Read, and J. Robbins. 2011. Underestimating the damage: interpreting cetacean carcass recoveries in the context of the Deepwater

Horizon/BP incident. Conserv. Letters 4:228-233.

Wilson K, Littnan C, Read AJ. 2017. Movements and home ranges of monk seals in the main Hawaiian Islands.

Marine Mammal Science. 33:1080-96.